

Geometry & Topology Monographs

Volume 7: Proceedings of the Casson Fest

Pages 135–144

On the additivity of knot width

MARTIN SCHARLEMANN

ABIGAIL THOMPSON

Abstract It has been conjectured that the geometric invariant of knots in 3-space called the *width* is nearly additive. That is, letting $w(K) \in 2\mathbb{N}$ denote the width of a knot $K \subset S^3$, the conjecture is that $w(K \# K') = w(K) + w(K') - 2$. We give an example of a knot K_1 so that for K_2 any 2-bridge knot, it appears that $w(K_1 \# K_2) = w(K_1)$, contradicting the conjecture.

AMS Classification 11Y16, 57M50; 57M25

Keywords Knot, width, additivity, Haken surfaces

Dedicated to Andrew Casson, a mathematician's mathematician

1 Background

In [1] Gabai associated to a knot in 3-space an even number called its width (see Definition 2 below for the precise definition). Width can be viewed as a generalization of bridge number, an invariant first studied by Schubert [4]. Schubert's remarkable discovery was that, for $b(K)$ the bridge number of $K \subset S^3$, it is always true that

$$b(K \# K') = b(K) + b(K') - 1.$$

An alternative formulation of what Schubert showed is this: one way of putting the knot sum $K \# K'$ into minimal bridge position is to put both K, K' into minimal bridge position, then place them vertically adjacent in S^3 . Then create their knot sum via a vertical band (see Figure 1).

Because width generalizes the notion of bridge number, it is natural to hope that a similar equality is true for width. Just as the construction described above and shown in Figure 1 makes it obvious that

$$b(K \# K') \leq b(K) + b(K') - 1$$

(Schubert's deep contribution is the proof of the reverse inequality, see [5]), it also shows that knot width w satisfies the inequality

$$w(K \# K') \leq w(K) + w(K') - 2.$$

That is, the resulting presentation of $K \# K'$ has width precisely equal to $w(K) + w(K') - 2$. The inequality reflects uncertainty over whether this presentation is of minimal width (that is, *thin*) among all possible presentations for $K \# K'$. This suggests the following

Conjecture 1 For all knots $K, K' \subset S^3$,

$$w(K \# K') = w(K) + w(K') - 2.$$




Figure 1

Beyond the analogy with bridge number, and the ease of the inequality in one direction, there are two pieces of evidence for Conjecture 1. One piece of evidence is the proof in [2] that the conjecture is true for knots that are *small*, that is, knots which have no closed essential surfaces in their complements. A second but weaker piece of evidence is the proof in [3] that in any case,

$$w(K \# K') \geq \max\{w(K), w(K')\}.$$

This shows that at least some inequality in the reverse direction is true.

The aim here is to present an example of a knot (actually a family of knots) K_1 which appears to have the property that

$$w(K_1 \# K_2) = w(K_1)$$

whenever K_2 is a 2-bridge knot. Since the width of a 2-bridge knot is 8, this would be a counterexample to Conjecture 1. Although we are not able to prove this equality, the construction of K_1 is so flexible that the example does seem to undercut any hope that the conjecture is true. Only the absence so far of a good method to prove that our presentation of the knot (family) K_1 is the presentation of least width stands in the way of a complete proof that at least some of these knots are counterexamples to Conjecture 1.

2 The example

The example K_1 is shown in Figure 2. It is actually a family of examples, because specific braids inside of boxes are not specified; the point will be that, in search of an example, the number of critical points r can be made arbitrarily high and the braids themselves be made arbitrarily complicated.




Figure 2

Figure 3 shows an imbedding of $K_1 \# (\text{trefoil})$ into S^3 ; a decomposing sphere

is shown in the figure. Note that the width of the presentations of K_1 and $K_1 \# (\text{trefoil})$ with respect to the vertical height function are the same, since there is an obvious level-preserving reimbedding of one to the other. (The construction of K_1 was inspired by the extensive use of level-preserving reimbeddings in [3].) Moreover, the only property of the trefoil knot that is used in the level-preserving reimbedding is the fact that it is 2-bridge — any other 2-bridge knot would do.




Figure 3

If we knew that the given presentation of the knot K_1 is thin (that is, of minimal width) we would be done. We do not yet know a full proof of this, but cite two pieces of evidence:

First of all, the braids in the four braid boxes of Figure 2 may be made very complicated so that the number of critical points of a plane sweeping across the boxes in any direction except the vertical will have a large number of critical points and arbitrarily high width. Although this does not obviously rule out the possibility of more complicated sweep-outs, it seems very likely that with sufficiently complicated braids in place a sweep-out can only be thin for the whole knot if the sweep-out passes through these braid-boxes vertically, that is as horizontal planes, just as in the given presentation. (This is a stronger requirement than merely ensuring that the tangles within the braid boxes are

thin.) Once this is ensured, the order in which the sweep-out passes through each tangle could be compared to the given order; alternative orders, such as the order shown in Figure 4, will give rise to greater width (see calculation below) as long as $r \geq 4$.

This second but indirect bit of evidence that K_1 is thin is this: According to Schubert's theorem, there is a presentation of K_1 with lower bridge number than that of $K_1 \# (\text{trefoil})$ shown in Figure 3. But the Figure 3 presentation has the same bridge number as the presentation of K_1 shown in Figure 2. Given that width is a generalization of bridge number, we ought to check that the presentation that lowers bridge number from that in Figure 2 does not also lower width. Figure 4 shows a presentation of K_1 that has lower bridge number than the original presentation, shown in Figure 2. But in fact the presentation of K_1 in Figure 4 is not as thin as that in Figure 2.




Figure 4

To explicitly compare the widths of the two presentations, recall the definitions (see [3]):

Definition 2 Let $p: S^3 \rightarrow \mathbb{R}$ be the standard height function and for each $-1 < t < 1$ let S^t denote the sphere $p^{-1}(t)$. Let $K \subset S^3$ be a knot in general position with respect to p and $c_1, \dots, c_n$ be the critical values of $p|_K$ listed in increasing order $c_1 < \dots < c_n$. Choose $r_1, \dots, r_{n-1}$ so that $c_i < r_i < c_{i+1}$ for $i = 1, \dots, n-1$. The *width of K with respect to p* , denoted by $w(K, p)$, is $\sum_i |K \cap S^{r_i}|$. The *width of K* , denoted by $w(K)$, is the minimum of $w(K', p)$ over all knots K' isotopic to K . We say that K is in *thin position* if $w(K, p) = w(K)$.

The key operational feature of width is that it is reduced if a minimum is pushed up above a maximum whereas switching the levels of two adjacent minima or two adjacent maxima has no effect.

There is an easier way to calculate width. For the levels r_i described above, call r_i a *thin level* of K (and the sphere S^{r_i} a *thin sphere* for K) with respect to p if c_i is a maximum value for $p|_K$ and c_{i+1} is a minimum value for $p|_K$. Dually r_i is a *thick level* of K (and the sphere S^{r_i} a *thick sphere* for K) with respect to p if c_i is a minimum value for $p|_K$ and c_{i+1} is a maximum value for $p|_K$. Since the lowest critical point of $p|_K$ is a minimum and the highest is a maximum, there is one more thick level than thin level.

Lemma 3 Let $r_{i_1}, \dots, r_{i_k}$ be the thick levels of K and $r_{j_1}, \dots, r_{j_{k-1}}$ the thin levels. Set $a_l = |K \cap S^{r_{i_l}}|$ and $b_l = |K \cap S^{r_{j_l}}|$. Then

$$2w(K) = \sum_{l=1}^k a_l^2 - \sum_{l=1}^{k-1} b_l^2.$$

Proof See [3]. □

Now apply this formula to the presentations of K_1 given in Figures 2 and 4, with thick and thin spheres (appearing as planes) noted in the figures. The former has width

$$4(r+1)^2 + 50 - 16 = 2(2r^2 + 4r + 19).$$

The latter has width

$$4(r+2)^2 - 8 = 2(2r^2 + 8r + 4).$$

So as long as $r \geq 4 \Rightarrow 8r + 4 > 4r + 19$ the presentation of K_1 in Figure 2 is thinner.

3 Higher bridge number

It is reasonable to ask whether likely counterexamples to the conjecture are limited to knots, such as those above, whose width is unchanged by adding 2-bridge knots. After all, 2-bridge knots often play a special role in knot theory. Almost certainly the answer is no; in this section we briefly note how a 3-bridge counterexample might be constructed. In principle the same ideas should work with arbitrary bridge number, though this would appear to be increasingly difficult to demonstrate.




Figure 5

Consider a knot K_3 of the form described in Figure 5. The notation r_i denotes the number of strands of the part of the knot that lies inside the “braid-box”, that is, twice the number of maxima appearing above the upper boxes (respectively, below the lower boxes). The notation s_i refers to the number of parallel strands between the boxes. The thin unmarked strand represents a single strand. We have earlier seen that the knot K_1 in Figure 2 can be added to any 2-bridge knot, without apparently affecting its width, by reimbedding it in a level-preserving way as a satellite of any 2-bridge knot. In the same way, the knot K_3 in Figure 5 can be reimbedded in a level-preserving way as a satellite of any 3-bridge knot K_4 . This reimbedding gives a presentation of the sum $K_3 \# K_4$ that has the same width as the original presentation of K_3 . So if the given presentation of K_3 is thin, we have a counterexample to Conjecture 1 in which one of the summand knots is the specific knot K_3 and the other is

any 3-bridge knot.

Of course there is still the difficulty of showing that the presentation of K_3 in Figure 5 is thin. But again the flexibility in how the braid-boxes are filled in suggests that filling in with complicated braids will force any thin presentation also to sweep across the braid boxes vertically. So a thin presentation is likely to preserve the braid-box structure. It is perhaps less plausible here than in the case of K_1 that a presentation which preserves the braid-box structure cannot be thinner than the original presentation; indeed, an example of a significantly different presentation of K_3 that preserves the braid-box structure is given in Figure 6.




Figure 6

However, it is straightforward (though not particularly easy) to show that with appropriate choice of r_i, s_j the first presentation is thinner than the second. A comparison of the two widths is unaffected by the size of r_3 . In addition, so long as we

- take the other values of r_i, s_j sufficiently large
- set r_1 only minimally larger than $s_1 + s_2$
- set $s_1 > s_3$, and, with these numbers set,
- take $r_2 > s_2 + s_3$ sufficiently high,

then the second presentation of K_3 will be wider than the first.

Another typical braid-box preserving presentation of K_3 is given in Figure 7. For this presentation, width can be made arbitrarily high simply by increasing r_3 sufficiently. As already noted, this has no effect on the comparison above, so ultimately we have values for r_i, s_j that guarantee the first presentation is thinnest of the three. Other braid-box preserving presentations are easily dealt with in a similar manner. As was the case for the 2-bridge examples of the previous section, there remains the challenge of showing that putting sufficiently complicated braids into the braid-boxes will guarantee that a thin presentation will preserve those braid-boxes.




Figure 7

Acknowledgement

This research was partially supported by NSF grants.

References

- [1] **D Gabai**, *Foliations and the topology of 3-manifolds. III*, J. Differential Geom. 26 (1987) 479–536 MR0910018
- [2] **Y Rieck**, **E Sedgwick**, *Thin position for a connected sum of small knots*, Algebr. Geom. Topol. 2 (2002) 297–309 MR1917054

- [3] **M Scharlemann, J Schultens**, *3-manifolds with planar presentations and the width of satellite knots*, [arXiv:math.GT/0304271](#)
- [4] **H Schubert**, *Über eine numerische Knoteninvariante*, Math. Z. 61 (1954) 245–288 MR0072483
- [5] **J Schultens**, *Additivity of bridge numbers of knots*, Math. Proc. Cambridge Philos. Soc. 135 (2003) 539–544 MR2018265

*Mathematics Department, University of California
Santa Barbara, CA 93106, USA*

and

*Mathematics Department, University of California
Davis, CA 95616, USA*

Email: mgscharl@math.ucsb.edu, thompson@math.ucdavis.edu

Received: 19 March 2004 Revised: 28 July 2004